

## **WO04032937**

Publication Title:

WO04032937

Abstract:

Abstract not available for WO04032937 Data supplied from the esp@cenet database - Worldwide

-----  
Courtesy of <http://v3.espacenet.com>

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
22 April 2004 (22.04.2004)

PCT

(10) International Publication Number  
**WO 2004/032937 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 31/517**, 31/7068, A61P 35/00
- (21) International Application Number:  
PCT/GB2003/004334
- (22) International Filing Date: 6 October 2003 (06.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
0223380.7 9 October 2002 (09.10.2002) GB
- (71) Applicant (*for all designated States except MG, US*): **ASTRAZENECA AB** [SE/SE]; S-151 85 Sodertalje (SE).
- (71) Applicant (*for MG only*): **ASTRAZENECA UK LIMITED** [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **BARGE, Alan** [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
- (74) Agent: **ASTRAZENECA**; Global Intellectual Property, P O Box 272, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
- with international search report
  - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF THE QUINAZOLINE DERIVATIVE ZD6474 COMBINED WITH GEMCITABINE AND OPTIONALLY IONISING RADIATION IN THE TREATMENT OF DISEASES ASSOCIATED WITH ANGIOGENESIS AND/OR INCREASED VASCULAR PERMEABILITY

(57) Abstract: The present invention relates to a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation, particularly a method for the treatment of a cancer, particularly a cancer involving a solid tumour, which comprises the administration of ZD6474 in combination with gemcitabine; to a pharmaceutical composition comprising ZD6474 and gemcitabine; to a combination product comprising ZD6474 and gemcitabine for use in a method of treatment of a human or animal body by therapy; to a kit comprising ZD6474 and gemcitabine; to the use of ZD6474 and gemcitabine in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation.

WO 2004/032937 A1

USE OF THE QUINAZOLINE DERIVATIVE ZD6474 COMBINED WITH GEMCITABINE AND  
OPTIONALLY IONISING RADIATION IN THE TREATMENT OF DISEASES ASSOCIATED  
WITH ANGIOGENESIS AND/OR INCREASED VASCULAR PERMEABILITY

The present invention relates to a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which  
5 is optionally being treated with ionising radiation, particularly a method for the treatment of a cancer, particularly a cancer involving a solid tumour, which comprises the administration of ZD6474 in combination with gemcitabine; to a pharmaceutical composition comprising ZD6474 and gemcitabine; to a combination product comprising ZD6474 and gemcitabine for use in a method of treatment of a human or animal body by therapy; to a kit comprising  
10 ZD6474 and gemcitabine; to the use of ZD6474 and gemcitabine in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation.

Normal angiogenesis plays an important role in a variety of processes including  
15 embryonic development, wound healing and several components of female reproductive function. Undesirable or pathological angiogenesis has been associated with disease states including diabetic retinopathy, psoriasis, cancer, rheumatoid arthritis, atheroma, Kaposi's sarcoma and haemangioma (Fan et al, 1995, Trends Pharmacol. Sci. 16: 57-66; Folkman, 1995, Nature Medicine 1: 27-31). Alteration of vascular permeability is thought to play a role in both  
20 normal and pathological physiological processes (Cullinan-Bove et al, 1993, Endocrinology 133: 829-837; Senger et al, 1993, Cancer and Metastasis Reviews, 12: 303-324). Several polypeptides with *in vitro* endothelial cell growth promoting activity have been identified including, acidic and basic fibroblast growth factors (aFGF & bFGF) and vascular endothelial growth factor (VEGF). By virtue of the restricted expression of its receptors, the growth factor  
25 activity of VEGF, in contrast to that of the FGFs, is relatively specific towards endothelial cells. Recent evidence indicates that VEGF is an important stimulator of both normal and pathological angiogenesis (Jakeman et al, 1993, Endocrinology, 133: 848-859; Kolch et al, 1995, Breast Cancer Research and Treatment, 36:139-155) and vascular permeability (Connolly et al, 1989, J. Biol. Chem. 264: 20017-20024). Antagonism of VEGF action by sequestration of VEGF with  
30 antibody can result in inhibition of tumour growth (Kim et al, 1993, Nature 362: 841-844).

Receptor tyrosine kinases (RTKs) are important in the transmission of biochemical signals across the plasma membrane of cells. These transmembrane molecules characteristically

consist of an extracellular ligand-binding domain connected through a segment in the plasma membrane to an intracellular tyrosine kinase domain. Binding of ligand to the receptor results in stimulation of the receptor-associated tyrosine kinase activity which leads to phosphorylation of tyrosine residues on both the receptor and other intracellular molecules. These changes in

5 tyrosine phosphorylation initiate a signalling cascade leading to a variety of cellular responses. To date, at least nineteen distinct RTK subfamilies, defined by amino acid sequence homology, have been identified. One of these subfamilies is presently comprised by the *fms*-like tyrosine kinase receptor, Flt-1, the kinase insert domain-containing receptor, KDR (also referred to as Flk-1), and another *fms*-like tyrosine kinase receptor, Flt-4. Two of these related RTKs, Flt-1


10 and KDR, have been shown to bind VEGF with high affinity (De Vries et al, 1992, Science 255: 989-991; Terman et al, 1992, Biochem. Biophys. Res. Comm. 1992, 187: 1579-1586). Binding of VEGF to these receptors expressed in heterologous cells has been associated with changes in the tyrosine phosphorylation status of cellular proteins and calcium fluxes.

Quinazoline derivatives which are inhibitors of VEGF receptor tyrosine kinase are

15 described in International Patent Applications Publication Nos. WO 98/13354 and WO 01/32651. In WO 98/13354 and WO 01/32651 compounds are described which possess activity against VEGF receptor tyrosine kinase (VEGF RTK) whilst possessing some activity against epidermal growth factor (EGF) receptor tyrosine kinase (EGF RTK).

ZD6474 is 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline:

20



ZD6474

25 ZD6474 falls within the broad general disclosure of WO 98/13354 and is exemplified in WO 01/32651. ZD6474 is a potent inhibitor of VEGF RTK and also has some activity against EGF RTK. ZD6474 has been shown to elicit broad-spectrum anti-tumour activity in a range

of models following once-daily oral administration (Wedge S.R., Ogilvie D.J., Dukes M. et al, Proc. Am. Assoc. Canc. Res. 2001; 42: abstract 3126).

In WO 98/13354 and WO 01/32651 it is stated that compounds of their inventions:  
“may be applied as a sole therapy or may involve, in addition to a compound of the invention,  
5 one or more other substances and/or treatments. Such conjoint treatment may be achieved by  
way of the simultaneous, sequential or separate administration of the individual components of  
the treatment.”

WO 98/13354 and WO 01/32651 then go on to describe examples of such conjoint treatment  
including surgery, radiotherapy and various types of chemotherapeutic agent including inhibitors  
10 of growth factor function.

Nowhere in WO 98/13354 and WO 01/32651 does it suggest the combination of a  
compound of the invention and gemcitabine for the treatment of any disease state including  
cancer.

Nowhere in WO 98/13354 and WO 01/32651 is the specific combination of ZD6474  
15 and gemcitabine suggested.

Nowhere in WO 98/13354 and WO 01/32651 does it state that use of any compound  
of the invention therein with other treatments will produce surprisingly beneficial effects.

A triple combination of a VEGF RTK inhibitor (PTK 787), an EGF RTK inhibitor (PKI  
166) and gemcitabine is described in Baker et al, Cancer Research 62, 1996 2003, April 1,  
20 2002. The authors concluded that a combination of either PTK 787 with gemcitabine or PKI  
166 with gemcitabine was beneficial but that the triple combination did not produce additive  
therapeutic effects.

A combination of gemcitabine with DC101, a VEGF receptor-2 antibody (anti-KDR  
antibody) is described in Bruns et al, International Journal of Cancer vol 102, issue 2, 2002  
25 pages 101-108.

Unexpectedly and surprisingly we have now found that the particular compound  
ZD6474 used in combination with a particular selection from the broad description of  
combination therapies listed in WO 98/13354 and WO 01/32651, namely with gemcitabine,  
produces significantly better effects than any one of ZD6474 and gemcitabine used alone. In  
30 particular, ZD6474 used in combination with gemcitabine produces significantly better effects  
on solid tumours than any one of ZD6474 and gemcitabine used alone.

Gemcitabine is (INN) 2'-deoxy-2',2'-difluorocytidine monohydrochloride ( $\beta$ -isomer).

Gemcitabine is also known as Gemzar <sup>TM</sup> (Trademark of Lilly) and it is a cytotoxic agent. It is an anti-metabolite which causes inhibition of DNA synthesis.

Anti-cancer effects of a method of treatment of the present invention include, but are not limited to, anti-tumour effects, the response rate, the time to disease progression and the survival rate. Anti-tumour effects of a method of treatment of the present invention include but are not limited to, inhibition of tumour growth, tumour growth delay, regression of tumour, shrinkage of tumour, increased time to regrowth of tumour on cessation of treatment, slowing of disease progression. It is expected that when a method of treatment of the present invention is administered to a warm-blooded animal such as a human, in need of treatment for cancer, with or without a solid tumour, said method of treatment will produce an effect, as measured by, for example, one or more of: the extent of the anti-tumour effect, the response rate, the time to disease progression and the survival rate.

According to the present invention there is provided a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine.

According to a further aspect of the present invention there is provided a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine; wherein ZD6474 and gemcitabine

may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises  
5 administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine; wherein ZD6474 and gemcitabine may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for  
10 the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine; wherein ZD6474 and gemcitabine may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

15 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises ZD6474 or a pharmaceutically acceptable salt thereof, and gemcitabine, in association with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a combination product comprising ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine, for  
20 use in a method of treatment of a human or animal body by therapy.

According to a further aspect of the present invention there is provided a kit comprising ZD6474 or a pharmaceutically acceptable salt thereof, and gemcitabine.

According to a further aspect of the present invention there is provided a kit comprising:

- 25 a) ZD6474 or a pharmaceutically acceptable salt thereof in a first unit dosage form;  
b) gemcitabine in a second unit dosage form; and  
c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- 30 a) ZD6474 or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable excipient or carrier, in a first unit dosage form;

b) gemcitabine together with a pharmaceutically acceptable excipient or carrier, in a second unit dosage form; and

c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided the use of  
5 ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human.

According to a further aspect of the present invention there is provided the use of  
ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a  
10 medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as a human.

According to a further aspect of the present invention there is provided the use of  
ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a  
medicament for use in the production of an anti-tumour effect in a warm-blooded animal such  
15 as a human.

According to a further aspect of the present invention there is provided a therapeutic combination treatment comprising the administration of an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, optionally together with a pharmaceutically acceptable excipient or carrier, and the simultaneous, sequential or separate administration of  
20 an effective amount of gemcitabine, wherein gemcitabine may optionally be administered together with a pharmaceutically acceptable excipient or carrier, to a warm-blooded animal such as a human in need of such therapeutic treatment.

Such therapeutic treatment includes an antiangiogenic and/or vascular permeability effect, an anti-cancer effect and an anti-tumour effect.

25 A combination treatment of the present invention as defined herein may be achieved by way of the simultaneous, sequential or separate administration of the individual components of said treatment. A combination treatment as defined herein may be applied as a sole therapy or may involve surgery or radiotherapy or an additional chemotherapeutic agent in addition to a combination treatment of the invention.

30 Surgery may comprise the step of partial or complete tumour resection, prior to, during or after the administration of the combination treatment with ZD6474 described herein.



Other chemotherapeutic agents for optional use with a combination treatment of the present invention include those described in WO 01/32651 which is incorporated herein by reference. Such chemotherapy may cover five main categories of therapeutic agent:

- (i) other antiangiogenic agents including vascular targeting agents;
- 5 (ii) cytostatic agents;
- (iii) biological response modifiers (for example interferon);
- (iv) antibodies (for example edrecolomab); and
- (v) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology.

- 10 A particular example of a chemotherapeutic agent for use with a combination treatment of the present invention is a platinum anti-cancer agent such as cisplatin; such a combination is expected to be particularly useful for the treatment of lung cancer and bladder cancer.

The administration of a triple combination of ZD6474, gemcitabine and ionising radiation may produce effects, such as anti-tumour effects, greater than those achieved with  
15 any of ZD6474, gemcitabine and ionising radiation used alone, greater than those achieved with the combination of ZD6474 and gemcitabine, greater than those achieved with the combination of ZD6474 and ionising radiation, greater than those achieved with the combination of gemcitabine and ionising radiation.

According to the present invention there is provided a method for the production of an  
20 antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation.

25 According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation.

30 According to a further aspect of the present invention there is provided a method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a

pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation.

According to a further aspect of the present invention there is provided a method for  
5 the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation, wherein ZD6474 and gemcitabine may each  
10 optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and  
15 before, after or simultaneously with an effective amount of ionising radiation, wherein ZD6474 and gemcitabine may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human,  
20 which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation, wherein ZD6474 and gemcitabine may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

25 According to a further aspect of the present invention there is provided the use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

30 According to a further aspect of the present invention there is provided the use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a

medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

According to a further aspect of the present invention there is provided the use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a  
5 medicament for use in the production of an anti-tumour effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

According to a further aspect of the present invention there is provided a therapeutic combination treatment comprising the administration of an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, optionally together with a pharmaceutically  
10 acceptable excipient or carrier, and the administration of an effective amount of gemcitabine, optionally together with a pharmaceutically acceptable excipient or carrier and the administration of an effective amount of ionising radiation, to a warm-blooded animal such as a human in need of such therapeutic treatment wherein the ZD6474, gemcitabine and ionising radiation may be administered simultaneously, sequentially or separately and in any order.

15 A warm-blooded animal such as a human which is being treated with ionising radiation means a warm-blooded animal such as a human which is treated with ionising radiation before, after or at the same time as the administration of a medicament or combination treatment comprising ZD6474 and gemcitabine. For example said ionising radiation may be given to said warm-blooded animal such as a human within the period of a week before to a week after the  
20 administration of a medicament or combination treatment comprising ZD6474 and gemcitabine. This means that ZD6474, gemcitabine and ionising radiation may be administered separately or sequentially in any order, or may be administered simultaneously. The warm-blooded animal may experience the effect of each of ZD6474, gemcitabine and radiation simultaneously.

25 According to one aspect of the present invention the ionising radiation is administered before one of ZD6474 and gemcitabine or after one of ZD6474 and gemcitabine.

According to one aspect of the present invention the ionising radiation is administered before both ZD6474 and gemcitabine or after both ZD6474 and gemcitabine.

According to one aspect of the present invention ZD6474 is administered to a warm-  
30 blooded animal after the animal has been treated with ionising radiation.

According to another aspect of the present invention the effect of a method of treatment of the present invention is expected to be at least equivalent to the addition of the

effects of each of the components of said treatment used alone, that is, of each of ZD6474 and gemcitabine used alone or of each of ZD6474, gemcitabine and ionising radiation used alone.

According to another aspect of the present invention the effect of a method of treatment of the present invention is expected to be greater than the addition of the effects of  
5 each of the components of said treatment used alone, that is, of each of ZD6474 and gemcitabine used alone or of each of ZD6474, gemcitabine and ionising radiation used alone.

According to another aspect of the present invention the effect of a method of treatment of the present invention is expected to be a synergistic effect.

According to the present invention a combination treatment is defined as affording a  
10 synergistic effect if the effect is therapeutically superior, as measured by, for example, the extent of the response, the response rate, the time to disease progression or the survival period, to that achievable on dosing one or other of the components of the combination treatment at its conventional dose. For example, the effect of the combination treatment is synergistic if the effect is therapeutically superior to the effect achievable with ZD6474 or gemcitabine or  
15 ionising radiation alone. Further, the effect of the combination treatment is synergistic if a beneficial effect is obtained in a group of patients that does not respond (or responds poorly) to ZD6474 or gemcitabine or ionising radiation alone. In addition, the effect of the combination treatment is defined as affording a synergistic effect if one of the components is dosed at its conventional dose and the other component(s) is/are dosed at a reduced dose and  
20 the therapeutic effect, as measured by, for example, the extent of the response, the response rate, the time to disease progression or the survival period, is equivalent to that achievable on dosing conventional amounts of the components of the combination treatment. In particular, synergy is deemed to be present if the conventional dose of ZD6474 or gemcitabine or ionising radiation may be reduced without detriment to one or more of the extent of the response, the  
25 response rate, the time to disease progression and survival data, in particular without detriment to the duration of the response, but with fewer and/or less troublesome side-effects than those that occur when conventional doses of each component are used.

As stated above the combination treatments of the present invention as defined herein are of interest for their antiangiogenic and/or vascular permeability effects. Angiogenesis  
30 and/or an increase in vascular permeability is present in a wide range of disease states including cancer (including leukaemia, multiple myeloma and lymphoma), diabetes, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, arterial

restenosis, autoimmune diseases, acute inflammation, lymphoedema, endometriosis, dysfunctional uterine bleeding and ocular diseases with retinal vessel proliferation including age-related macular degeneration. Combination treatments of the present invention are expected to be particularly useful in the prophylaxis and treatment of diseases such as cancer  
5 and Kaposi's sarcoma. In particular such combination treatments of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, pancreas, bladder, breast, prostate, lungs and skin. More especially combination treatments of the present invention are expected to slow advantageously the growth of tumours in pancreatic cancer and lung cancer, for example mesothelioma and non-small cell  
10 lung cancer (NSCLC). More particularly such combination treatments of the invention are expected to inhibit any form of cancer associated with VEGF including leukaemia, multiple myeloma and lymphoma and also, for example, to inhibit the growth of those primary and recurrent solid tumours which are associated with VEGF, especially those tumours which are significantly dependent on VEGF for their growth and spread, including for example, certain  
15 tumours of the colon, pancreas, bladder, breast, prostate, lung, vulva and skin, particularly NSCLC.

In another aspect of the present invention ZD6474 and gemcitabine, optionally with ionising radiation, are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with VEGF especially those tumours which are significantly  
20 dependent on VEGF for their growth and spread.

In another aspect of the present invention ZD6474 and gemcitabine, optionally with ionising radiation, are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with both VEGF and EGF especially those tumours which are significantly dependent on VEGF and EGF for their growth and spread.

25 The compositions described herein may be in a form suitable for oral administration, for example as a tablet or capsule, for nasal administration or administration by inhalation, for example as a powder or solution, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) for example as a sterile solution, suspension or emulsion, for topical administration for example as an ointment or cream, for rectal  
30 administration for example as a suppository or the route of administration may be by direct injection into the tumour or by regional delivery or by local delivery. In other embodiments of the present invention the ZD6474 of the combination treatment may be delivered

endoscopically, intratracheally, intralesionally, percutaneously, intravenously, subcutaneously, intraperitoneally or intratumorally. Preferably ZD6474 is administered orally. In general the compositions described herein may be prepared in a conventional manner using conventional excipients. The compositions of the present invention are advantageously presented in unit  
5 dosage form.

ZD6474 will normally be administered to a warm-blooded animal at a unit dose within the range 10-500mg per square metre body area of the animal, for example approximately 0.3-15mg/kg in a human. A unit dose in the range, for example, 0.3-15mg/kg, preferably 0.5-5mg/kg is envisaged and this is normally a therapeutically-effective dose. A unit dosage  
10 form such as a tablet or capsule will usually contain, for example 25-500mg of active ingredient. Preferably a daily dose in the range of 0.5-5mg/kg is employed.

Gemcitabine may be administered according to known clinical practice. For example in NSCLC the recommended dose of gemcitabine is 1000mg/m<sup>2</sup> given by 30 minute intravenous infusion. This may be repeated once weekly for three weeks, followed by a one week rest  
15 period. This four week cycle may then be repeated. Dosage reduction may be necessary if the patient experiences undue toxicity. In pancreatic cancer the recommended dose of gemcitabine is 1000mg/m<sup>2</sup> given by 30 minute intravenous infusion. This may be repeated once weekly for seven weeks followed by a week of rest. Subsequent cycles may consist of injections once weekly for three consecutive weeks out of every four weeks. Dosage reduction may be  
20 necessary if the patient experiences undue toxicity.

The dosages and schedules may vary according to the particular disease state and the overall condition of the patient. Dosages and schedules may also vary if, in addition to a combination treatment of the present invention, one or more additional chemotherapeutic agents is/are used. Scheduling can be determined by the practitioner who is treating any particular patient.

25 Radiotherapy may be administered according to the known practices in clinical radiotherapy. The dosages of ionising radiation will be those known for use in clinical radiotherapy. The radiation therapy used will include for example the use of  $\gamma$ -rays, X-rays, and/or the directed delivery of radiation from radioisotopes. Other forms of DNA damaging factors are also included in the present invention such as microwaves and UV-irradiation. For  
30 example X-rays may be dosed in daily doses of 1.8-2.0Gy, 5 days a week for 5-6 weeks. Normally a total fractionated dose will lie in the range 45-60Gy. Single larger doses, for example 5-10Gy may be administered as part of a course of radiotherapy. Single doses may be

administered intraoperatively. Hyperfractionated radiotherapy may be used whereby small doses of X-rays are administered regularly over a period of time, for example 0.1Gy per hour over a number of days. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and on the uptake by cells.

5 As stated above the size of the dose of each therapy which is required for the therapeutic or prophylactic treatment of a particular disease state will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient. For example, it may be necessary or desirable to reduce the  
10 above-mentioned doses of the components of the combination treatments in order to reduce toxicity.

The present invention relates to combinations of gemcitabine with ZD6474 or with a salt of ZD6474.

Salts of ZD6474 for use in pharmaceutical compositions will be pharmaceutically  
15 acceptable salts, but other salts may be useful in the production of ZD6474 and its pharmaceutically acceptable salts. Such salts may be formed with an inorganic or organic base which affords a pharmaceutically acceptable cation. Such salts with inorganic or organic bases include for example an alkali metal salt, such as a sodium or potassium salt, an alkaline earth metal salt such as a calcium or magnesium salt, an ammonium salt or for example a salt with  
20 methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

ZD6474 may be synthesised according to any of the known processes for making ZD6474. For example ZD6474 may be made according to any of the processes described in WO 01/32651; for example those described in Examples 2(a), 2(b) and 2(c) of WO 01/32651.

25 Gemcitabine is commercially available.

The following tests may be used to demonstrate the activity of ZD6474 in combination with gemcitabine.

Model to investigate effects of ZD6474 and gemcitabine therapy on primary pancreatic tumour growth and metastases growing orthotopically in nude mice

30 Nude mice (n=10 per group) were injected with  $1 \times 10^6$  L3.6pl human pancreatic cancer cells into the pancreas. The mice received one of the following treatment regimes 8 days post-tumour injection:

Group 1 – control group received saline by daily oral application;

Group 2 - a single daily dose of ZD6474 (50mg/kg oral gavage);

Group 3 – gemcitabine twice weekly (on days 2 and 5 of each treatment week) (100mg/kg by intraperitoneal injection (i.p.));

5 Group 4 – a single daily dose of ZD6474 (50mg/kg oral gavage) and twice weekly gemcitabine (on days 2 and 5 of each treatment week) (GEM: 100mg/kg i.p.).

The animals were sacrificed 32 days after tumour cell injection.

The incidence of pancreatic tumours, liver metastases, lymph node metastases and peritoneal carcinosis was evaluated. The primary pancreatic tumour weight was measured. All

10 macroscopically enlarged lymph nodes and liver nodules were confirmed by histopathology (H&E staining).

The results are shown in Table 1

Table 1

Therapy Group	Incidence Pancreas	Liver Met	LN Met	Peritoneal Carcinosis	Tumour Weight Average +/- std dev	Body Weight Average +/- std dev
Control	10/10	6/10	10/10	3/10	mg 1231 +/- 290	g 21.4 +/- 2.2
ZD6474	10/10	1/10	3/10	1/10	541 +/- 201	19.2 +/- 2.9
Gemcitabine	9/9	4/9	9/9	3/9	836 +/- 291	21.7 +/- 2.1
ZD6474 + Gemcitabine	5/5	0/5	1/5	0/5	308 +/- 129	19.3 +/- 3.4

15

Compared with the average weight of control tumours (1231mg), tumours in treated animals reached an average weight of 836mg (gemcitabine) 541mg (ZD6474) and 308mg (gemcitabine + ZD6474). Liver metastases were present in 6/10 control and 4/9 gemcitabine treated animals, but only 1/10 and 0/5 animals on ZD6474 or combination treatment displayed liver

20 metastases, respectively. Lymph node metastases were present in 10/10 control and 9/9 gemcitabine treated animals, but only 3/10 and 1/5 animals on ZD6474 or combination treatment displayed lymph node metastases, respectively.



**CLAIMS**

1. A method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to  
5 said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine.
2. A method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a  
10 pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine.
3. A method according to claim 2 for the treatment of a cancer involving a solid tumour.
- 15 4. A pharmaceutical composition which comprises ZD6474 or a pharmaceutically acceptable salt thereof, and gemcitabine, in association with a pharmaceutically acceptable excipient or carrier.
5. A kit comprising ZD6474 or a pharmaceutically acceptable salt thereof, and  
20 gemcitabine.
6. Use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human.  
25
7. Use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as a human.
- 30 8. Use according to claim 7 wherein the anti-cancer effect comprises an anti-tumour effect.
9. A method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to

said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation.

- 5 10. A method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of gemcitabine and before, after or simultaneously with an effective amount of ionising radiation.

10

11. A method according to claim 10 for the treatment of a cancer involving a solid tumour.

12. Use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular  
15 permeability reducing effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

13. Use of ZD6474 or a pharmaceutically acceptable salt thereof and gemcitabine in the manufacture of a medicament for use in the production of an anti-cancer effect in a  
20 warm-blooded animal such as a human which is being treated with ionising radiation.

14. Use according to claim 13 wherein the anti-cancer effect comprises an anti-tumour effect.

## INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/GB 03/04334

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/517 A61K31/7068 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ, EMBASE, CHEM ABS Data, PASCAL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 01 32651 A (HENNEQUIN LAURENT FRANCOIS AND ;THOMAS ANDREW PETER (GB); ASTRAZEN)  10 May 2001 (2001-05-10)  cited in the application  page 1, line 3-8  page 2, line 16 -page 4, line 13  page 5, line 18  page 6, line 16-18  page 26, line 22 -page 28, line 17  ---  -/-</p>	1-14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

9 January 2004

Date of mailing of the international search report

13/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Houyvet, C

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 03/04334

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CIARDIELLO F ET AL: "Antitumor effects of ZD6474, a small molecule VEGF receptor tyrosine kinase inhibitor that is also active against EGF receptor tyrosine kinase"</p> <p>PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH ANNUAL, vol. 43, March 2002 (2002-03), pages 1080-1081, XP001157108</p> <p>93rd Annual Meeting of the American Association for Cancer Research; San Francisco, California, USA; April 06-10, 2002, March, 2002</p> <p>ISSN: 0197-016X</p> <p>abstract</p> <p>----</p>	1-14
A	<p>DREVS J ET AL: "Effect of ZD6474, a VEGF receptor tyrosine kinase inhibitor, on primary tumor growth, metastasis, vessel density and microvascular architecture in murine renal cell carcinoma"</p> <p>PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH ANNUAL, vol. 43, March 2002 (2002-03), page 1082</p> <p>XP001156998</p> <p>93rd Annual Meeting of the American Association for Cancer Research; San Francisco, California, USA; April 06-10, 2002, March, 2002</p> <p>ISSN: 0197-016X</p> <p>abstract</p> <p>----</p>	1-14
A	<p>WEDGE S R ET AL: "Combination of the VEGF receptor tyrosine kinase inhibitor ZD6474 and vascular-targeting agent ZD6126 produces an enhanced antitumor response"</p> <p>PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH ANNUAL, vol. 43, March 2002 (2002-03), page 1081</p> <p>XP001156999</p> <p>93rd Annual Meeting of the American Association for Cancer Research; San Francisco, California, USA; April 06-10, 2002, March, 2002</p> <p>ISSN: 0197-016X</p> <p>abstract</p> <p>----</p> <p>--- -/--</p>	1-14

## INTERNATIONAL SEARCH REPORT

 Int lional Application No  
 PCT/GB 03/04334

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>HOLDEN S ET AL: "Effects of ZD6474, an orally active inhibitor of VEGF receptor tyrosine kinase, in patients with solid tumors: Results from a phase I study" EUROPEAN JOURNAL OF CANCER, PERGAMON PRESS, OXFORD, GB, vol. 37, no. Supplement 6, October 2001 (2001-10), page S73 XP002256123 ISSN: 0959-8049 abstract</p> <p>----</p>	1-14
A	<p>HENNEQUIN L F ET AL: "Novel 4-anilinoquinazolines with C-7 basic side chains: Design and structure activity relationship of a series of potent, orally active, VEGF receptor tyrosine kinase inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 45, no. 6, 14 March 2002 (2002-03-14), pages 1300-1312, XP002256124 ISSN: 0022-2623 abstract; table 1</p> <p>----</p>	1-14
A	<p>GIACCONE G ET AL: "ZD1839 ('Iressa'), an orally-active, selective, epidermal growth factor receptor tyrosine kinase inhibitor (egfr-tki), is well tolerated in combination with gemcitabine and cisplatin, in patients with advanced solid tumours: Preliminary tolerability, efficacy and pharmacokinetic results" EUROPEAN JOURNAL OF CANCER, vol. 37, no. Supplement 6, October 2001 (2001-10), pages S30-S31, XP002266469 11th European Cancer Conference;Lisbon, Portugal; October 21-25, 2001 ISSN: 0959-8049 abstract</p> <p>-----</p>	1-14

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 03/04334

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0132651 A	10-05-2001	AU 1288601 A	14-05-2001
		BG 106659 A	31-03-2003
		BR 0015203 A	16-07-2002
		CA 2389767 A1	10-05-2001
		CN 1387527 T	25-12-2002
		CZ 20021526 A3	17-07-2002
		EE 200200237 A	16-06-2003
		EP 1244647 A1	02-10-2002
		WO 0132651 A1	10-05-2001
		HU 0203453 A2	28-05-2003
		JP 2003513089 T	08-04-2003
		NO 20022139 A	03-05-2002
		SK 6122002 A3	09-01-2003